

Product Datasheet

www.leinco.com



Anti-Mouse IL-4 (Clone 11B11) Purified *in vivo* GOLD™ Functional Grade Monoclonal Antibody

Product Information

Product No.: I-1071
Clone: 11B11
RRID: AB_2830405
Isotype: Rat IgG1 κ
Storage: Sterile 2-8°C

Product Description

Specificity:

Clone 11B11 recognizes an epitope on mouse IL-4.

Antigen Distribution:

IL-4 is secreted by Th2 cells, NKT cells, mast cells, eosinophils, and basophils.

Background:

IL-4 antibody, 11B11, recognizes mouse IL-4, also known as B-cell differentiation factor (BCDF) and B-cell stimulatory factor (BSF1). IL-4 is a pleiotropic 20 kDa cytokine secreted by T helper 2 cells (Th2) cells, natural killer T (NKT) cells, mast cells, eosinophils, and basophils^{1,2}. IL-4 signaling through the IL-4 receptor alpha chain (IL-4Ra) induces the differentiation of naive T cells to Th2 cells^{3,4} and stimulates B cell differentiation, proliferation, and class switching to IgG1 and IgE isotypes⁵. In addition, IL-4 is implicated in normal wound healing^{6,7} and airway inflammation in patients with asthma⁸.

Known Reactivity Species:

Mouse

Format:

in vivo GOLD™, Purified *in vivo* Functional Grade

Immunogen:

Partially Purified Native Mouse IL-4

Formulation

This monoclonal antibody is aseptically packaged and formulated in 0.01 M phosphate buffered saline (150 mM NaCl) PBS pH 7.2 - 7.4 with no carrier protein, potassium, calcium or preservatives added. Due to inherent biochemical properties of antibodies, certain products may be prone to precipitation over time. Precipitation may be removed by aseptic centrifugation and/or filtration.

Purity

≥95% monomer by analytical SEC, >95% by SDS Page

Endotoxin

< 1.0 EU/mg as determined by the LAL method

Storage and Stability

Functional grade preclinical antibodies may be stored sterile as received at 2-8°C for up to one month. For longer term storage, aseptically aliquot in working volumes without diluting and store at ≤ -70°C. **Avoid Repeated Freeze Thaw Cycles.**

Product Preparation

Functional grade preclinical antibodies are manufactured in an animal free facility using *in vitro* cell culture techniques and are purified by a multi-step process including the use of protein A or G to assure extremely low levels of endotoxins, leachable protein A or aggregates.

Applications

Applications and Recommended Usage (Quality Tested By Leinco):

Products are for research use only. Not for use in diagnostic or therapeutic procedures.

410 Axminster Dr, St. Louis, MO 63026

(800) 538-1145

leincoglobal@leinco.com

Product Datasheet

www.leinco.com



ELISA This antibody is useful as the capture antibody in a sandwich ELISA. The suggested coating concentration is 0.5-2.0 µg/ml. Titration of the reagent is recommended for optimal performance for each application.

Other Applications Reported in Literature:

N

IHC

ICC

IP

CyTOF®

FA

Country of Origin

USA

References

1. Yoshimoto T & Paul WE. (1994) J Exp Med. 179(4):1285–95
2. Kipnis J., et al. (2012) J Immunol. 189(9):4213–4219
3. Groth B., et al. (1992) J Exp Med. 176:1091–8
4. Murphy KM., et al. (1992) Proc Natl Acad Sci USA. 89:6065–9

Products are for research use only. Not for use in diagnostic or therapeutic procedures.

410 Axminster Dr, St. Louis, MO 63026

(800) 538-1145

leincoglobal@leinco.com