

# Mouse IgG<sub>2b</sub> Purified in vivo PLATINUM™ Functional Grade

## **Isotype Control**

**Product Information** 

Product No.: M1415 Clone: MPC-11

RRID: AB\_2831345 Isotype: Mouse IgG2b Storage: Sterile 2-8°C

## **Product Description**

## Specificity:

This Mouse Mouse isotype control antibody has been tested against selected species' cells and tissues to assure minimal cross reactivity.

#### Format:

Purified in vivo PLATINUM™ Functional Grade

#### **Formulation**

This monoclonal antibody is aseptically packaged and formulated in 0.01 M phosphate buffered saline (150 mM NaCl) PBS pH 7.2 - 7.4 with no carrier protein, potassium, calcium or preservatives added. Due to inherent biochemical properties of antibodies, certain products may be prone to precipitation over time. Precipitation may be removed by aseptic centrifugation and/or filtration.

#### **Purity**

≥98% monomer by analytical SEC, >95% by SDS Page

#### **Endotoxin**

< 0.5 EU/mg as determined by the LAL method

#### Storage and Stability

Functional grade preclinical antibodies may be stored sterile as received at 2-8°C for up to one month. For longer term storage, aseptically aliquot in working volumes without diluting and store at  $\leq$  -70°C.

#### Avoid Repeated Freeze Thaw Cycles.

#### **Product Preparation**

Functional grade preclinical antibodies are manufactured in an animal free facility using in vitro cell culture techniques and are purified by a multi-step process including the use of protein A or G to assure extremely low levels of endotoxins, leachable protein A or aggregates.

#### **Pathogen Testing**

To protect mouse colonies from infection by pathogens and to assure that experimental preclinical data is not affected by such pathogens, all of Leinco's Purified Functional PLATINUM™ antibodies are tested and guaranteed to be negative for all pathogens in the IDEXX IMPACT I Mouse Profile.

## **Product Datasheet**

#### www.leinco.com

### **Applications**

Applications and Recommended Usage (Quality Tested By Leinco):

**FC** This isotype control antibody should be used at the same concentration as the primary antibody.

Other Applications Reported in Literature:

FA

ICC

**ICFC** 

**IF Microscopy** 

**IF Staining** 

**WB** 

For specific conjugates of this clone, review literature for suggested application details.

## **Country of Origin**

USA