Leinco Technologies, Inc.
excellence in early discovery research™

Anti-Mouse I-A/I-E (MHC Class II)

Purified (PhenoCycler-Fusion (CODEX)® Ready)

Monoclonal Antibody

Product Information

Product No.:	H502
Clone:	M5/114.15.2
RRID:	AB_ 2893767
Isotype:	Rat IgG2b

Product Description

Specificity:
Clone M5/114.15.2 recognizes an epitope on the mouse MHC class II molecules I-A^b, I-A^d, I-A^q, I-E^d, and I-E^k. Clone M5/114.15.2 does not react with I-A^f, I-A^k, or I-A^s.

Antigen Distribution:
I-$A^{b/d/q}$ and I-$E^{d/k}$ are expressed on antigen-presenting cells, including macrophages, monocytes, DCs, and B cells, and activated T cells from mice of the I-$A^{b/d/q}$ and I-$E^{d/k}$ haplotypes.

Background:
MHC class II antibody, clone M5/114.15.2, recognizes the major histocompatibility complex (MHC) class II molecules I-A^b, I-A^d, I-A^q, I-E^d, and I-E^k. MHC class II is constitutively expressed on professional antigen-presenting cells (APCs), including macrophages/monocytes, dendritic cells (DCs), and B cells, and is induced on T cells upon activation[1]. MHC class II consists of two transmembrane proteins, a 35 kDa α (heavy) chain and 29 kDa β (light) chain[2]. The N-terminal α1 and β1 domains form the antigen-binding groove, which binds 13-25 aa peptides derived from exogenous antigens[3]. On APCs, MHC class II plays a critical role in the adaptive immune response by presenting phagocytosed antigens to helper CD4 T cells. The T cell receptor (TCR)/CD3 complex of CD4 T cells interacts with peptide-MHC class II, which induces CD4 T cell activation leading to the coordination and regulation of other effector cells. CD4 molecules also bind to MHC class II, which helps augment TCR signaling[4]. It has also been demonstrated that MHC class II express on activated T cells are capable of antigen presentation[5] and can transduce signals into T cells, enhancing T cell proliferation and activity[6].

Known Reactivity Species:
Mouse

Format:
Purified (PhenoCycler® Ready)

Immunogen:
C57BL/6 Spleen Cells

Formulation
This purified antibody is formulated in 0.01 M phosphate buffered saline (150 mM NaCl) PBS pH 7.4.

Products are for research use only. Not for use in diagnostic or therapeutic procedures.

410 Axminister Dr, St. Louis, MO 63026 (800) 538-1145 leincoglobal@leinco.com

Storage and Stability
This antibody is stable for at least one week when stored at 2-8°C. For long term storage, aliquot in working volumes without diluting and store at -20°C **in a manual defrost freezer.**
Avoid Repeated Freeze Thaw Cycles.

Applications
Applications and Recommended Usage (Quality Tested By Leinco):

CODEX® This I-A/I-E (Clone M5/114.15.2) antibody is formulated to simplify the antibody preparation needed when performing a CODEX® barcode conjugate. The suggested concentration is 0.5 mg/ml.

Country of Origin
USA

References
1. Holling TM, et al. (2004) Hum Immunol. 65(4):282-90
2. Mitaksov V & Fremont DH. (2006) J Biol Chem. 281(15):10618-25
3. Wieczorek M, et al. (2017) Front Immunol. 8:292
4. Artyomov MN, et al. (2010) Proc Natl Acad Sci USA. 107(39):16916-16921
5. Barnaba V, et al. (1994) Eur J Immunol. 24(1):71-5
6. Di Rosa F, et al. (1993) Hum Immunol. 38(4):251-60

Products are for research use only. Not for use in diagnostic or therapeutic procedures.
410 Axminister Dr, St. Louis, MO 63026 (800) 538-1145 leincoglobal@leinco.com