

## 8-Oxo-A DNA/RNA Damage Monoclonal Antibody

### ORDERING INFORMATION

Description	Anti-8-oxoA, a mouse monoclonal antibody targeting oxidative damage in RNA, DNA, ATP and other nucleotide precursors and degradation products
Catalog #:	12503
Sizes:	100 ug, 500 ug

### SPECIFICATION SUMMARY

Antigen Target:	8-Oxo-Adenosine
Specificity	This antibody recognizes 8-oxoA in single and double stranded RNA and DNA and also in free nucleotides such as 8-oxoATP
Background:	This target is a byproduct of oxidative damage to the adenosine nucleobase in DNA, RNA, and nucleotides such as ATP. In DNA, 8-oxoA mispairs may reduce fidelity of replication and cause transcriptional errors by RNA pol II. In RNA, 8-oxoA damage leads to attenuation of mRNA translation. Specialized base excision repair machinery and nucleotide sanitation processes are present to reduce the mutagenic consequences of 8-oxoA.
Host:	Mouse
Isotype:	IgG1κ
Immunogen:	8-Oxo-A conjugated to KLH
Clonality:	Monoclonal
Clone ID:	6E4
Target PubChem:	96852
Target Synonyms:	8-oxoA, 8-Oxoadenosine, 8-Hydroxyadenosine, 7,8-dihydro-8-oxoadenosine, 6-Azanyl-9-[(2r,3r,4s,5r)-3,4-Dihydroxy-5-(Hydroxymethyl)oxolan-2-Yl]-7h-Purin-8-One
Target Molecular Wt.	283.24 g/mol

### PROPERTIES

Form:	Liquid
Concentration:	Lot Specific
Formulation:	Phosphate Buffered Saline, pH 7.4
Cryo-preserved:	None
Anti-Microbial:	None
Purification:	Protein G Affinity Chromatography

*Products are for research use only. Not for use in diagnostic or therapeutic procedures.*

QED Bioscience, Inc. 410 Axminster Dr, St. Louis, MO 63026

Phone: (800) 538-1145

Email: [leincoglobal@leinco.com](mailto:leincoglobal@leinco.com)

## 8-Oxo-A DNA/RNA Damage Monoclonal Antibody


### APPLICATIONS

Applications:


ELISA

Dilution Instructions:

Dilute in PBS or medium that is identical to that used in the assay system. Optimal concentrations must be derived for specific applications.




Determined by a competitive inhibition ELISA with nucleoside variants. The %B/B0 is plotted as a function of the concentration (nM) of the indicated competitor. 100% B/B0 indicates no inhibition by the indicated competitor.



Selectivity and Specificity of 6E4 in RNA:

- \* Binds 8-oxoA in single and double stranded RNA
- \* Recognizes 8-oxoA in RNA base mispairs with standard geometry



Selectivity and Specificity of 6E4 in DNA:

- \* Binds 8-oxoA in single and double stranded DNA
- \* Recognizes 8-oxoA in all DNA base mispairs

### STORAGE AND STABILITY

Storage

Conditions:

Store antibody at -20C to -80C. Avoid multiple freeze-thaw cycles.

Stability:

Antibody is stable for at least 1 year when stored at -20C to -80C.

### REFERENCES

Daniel J. Tew a, Jakob M. Hebert b, Brad J. Schmier, Discovery and properties of a monoclonal antibody targeting 8-oxoA, an oxidized adenine lesion in DNA and RNA, Redox Biology 62 (2023) 102658, PMC10074937

Products are for research use only. Not for use in diagnostic or therapeutic procedures.

QED Bioscience, Inc. 410 Axminister Dr, St. Louis, MO 63026

Phone: (800) 538-1145

Email: leincoglobal@leincog.com